



## Original Article

## ECFS standards of care on CFTR-related disorders: Identification and care of the disorders



N.J. Simmonds<sup>a,b,\*</sup>, K.W. Southern<sup>c</sup>, E. De Wachter<sup>d</sup>, K. De Boeck<sup>e</sup>, F. Bodewes<sup>f</sup>, J.G. Mainz<sup>g</sup>, P.G. Middleton<sup>h</sup>, C. Schwarz<sup>i</sup>, V. Vloeberghs<sup>j</sup>, M. Wilschanski<sup>k</sup>, E. Bourrat<sup>l</sup>, J.D. Chalmers<sup>m</sup>, C.Y. Ooi<sup>n</sup>, D. Debray<sup>o</sup>, D.G. Downey<sup>p</sup>, P. Eschenhagen<sup>q</sup>, E. Girodon<sup>r</sup>, G. Hickman<sup>l</sup>, A. Koitschev<sup>s</sup>, D. Nazareth<sup>t</sup>, J.A. Nick<sup>u</sup>, D. Peckham<sup>v</sup>, D. VanDevanter<sup>w</sup>, C. Raynal<sup>x</sup>, I. Scheers<sup>y</sup>, M.D. Waller<sup>z</sup>, I. Sermet-Gaudelus<sup>aa</sup>, C. Castellani<sup>ab</sup>, on behalf of the ECFS Diagnostic Network Working Group

<sup>a</sup> Adult Cystic Fibrosis Centre, Royal Brompton Hospital, London, UK

<sup>b</sup> National Heart and Lung Institute, Imperial College London, UK

<sup>c</sup> Department of Women's and Children's Health, University of Liverpool, Alder Hey Children's Hospital, Liverpool, UK

<sup>d</sup> Cystic Fibrosis Center, Pediatric Pulmonology department, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel, Brussels, Belgium

<sup>e</sup> Department of Pediatrics, University of Leuven, Leuven, Belgium

<sup>f</sup> Pediatric Gastroenterology and Hepatology, Department of Pediatrics, University of Groningen Medical Center, Groningen, the Netherlands

<sup>g</sup> Cystic Fibrosis Center, Brandenburg Medical School (MHB), University, Klinikum Westbrandenburg, Brandenburg an der Havel, Germany

<sup>h</sup> Cystic Fibrosis and Bronchiectasis Service, Department of Respiratory and Sleep Medicine, Westmead Hospital, Sydney, New South Wales, Australia

<sup>i</sup> HMU-Health and Medical University Potsdam, CF Center Westbrandenburg, Campus Potsdam, Germany

<sup>j</sup> Brussels IVF, Centre for Reproductive Medicine, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel, Brussels, Belgium

<sup>k</sup> CF Center, Department of Pediatrics, Hadassah Medical Center and Faculty of Medicine, Hebrew University of Jerusalem, Jerusalem, Israel

<sup>l</sup> APHP, Service de Dermatologie, CRMR MAGEC Nord St Louis, Hôpital-Saint Louis, Paris, France

<sup>m</sup> Division of Molecular and Clinical Medicine, University of Dundee, Dundee, UK

<sup>n</sup> a) School of Clinical Medicine, Discipline of Paediatrics and Child Health, Medicine & Health, University of New South Wales, Level 8, Centre for Child Health Research & Innovation Bright Alliance Building Cnr Avoca & High Streets, Randwick, Sydney, NSW, Australia, 2031; b) Sydney Children's Hospital, Gastroenterology Department, High Street, Randwick, Sydney, NSW, Australia, 2031

<sup>o</sup> Pediatric Hepatology unit, Centre de Référence Maladies Rares (CRMR) de l'atrésie des voies biliaires et cholestases génétiques (AVB-CG), National network for rare liver diseases (Filfoie), ERN rare liver, Hôpital Necker-Enfants Malades, AP-HP, Université de Paris, Paris, France; Sorbonne Université, INSERM, Centre de Recherche Saint-Antoine (CRSA), Institute of Cardiometa-bolism and Nutrition (ICAN), Paris, France

<sup>p</sup> Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast, UK

<sup>q</sup> CF Center Westbrandenburg, Potsdam, Germany

<sup>r</sup> Service de Médecine Génomique des Maladies de Système et d'Organe, APHP.Centre - Université de Paris Cité, Hôpital Cochin, Paris, France

<sup>s</sup> Klinikum Stuttgart, Pediatric Otorhinolaryngology, Stuttgart, Germany

<sup>t</sup> a) Adult CF Unit, Liverpool Heart and Chest Hospital NHS Foundation Trust, U.K.; b) Clinical Infection, Microbiology and Immunology, University of Liverpool, UK

<sup>u</sup> Department of Medicine, National Jewish Health, Denver, CO, 80206, USA, Department of Medicine, University of Colorado School of Medicine, Aurora, CO, 80045, USA

<sup>v</sup> Leeds Institute of Medical Research, University of Leeds, Leeds, United Kingdom

<sup>w</sup> Department of Pediatrics, School of Medicine, Case Western Reserve University, Cleveland, Ohio, USA

<sup>x</sup> Laboratory of molecular genetics, University Hospital of Montpellier and INSERM U1046 PHYMEDEXP, Montpellier, France

<sup>y</sup> Department of Pediatrics, Pediatric Gastroenterology and Hepatology Unit, Cliniques Universitaires Saint Luc, Université Catholique de Louvain, Brussels, Belgium

<sup>z</sup> Adult Cystic Fibrosis and Respiratory Medicine, King's College Hospital NHS Foundation Trust, London, United Kingdom; Honorary Senior Lecturer, King's College London, London, United Kingdom

<sup>aa</sup> INSERM U1151, Institut Necker Enfants Malades, Paris, France; Université de Paris, Paris, France; Centre de référence Maladies Rares, Mucoviscidose et maladies apparentées, Hôpital Necker Enfants malades, Paris, France

<sup>ab</sup> IRCCS Istituto Giannina Gaslini, Cystic Fibrosis Center, Genoa, Italy

\* Corresponding author at: Adult Cystic Fibrosis Centre, Roycal Brompton Hospital, London, SW3 6NP, United Kingdom

E-mail address: [n.simmonds@imperial.ac.uk](mailto:n.simmonds@imperial.ac.uk) (N.J. Simmonds).

<https://doi.org/10.1016/j.jcf.2024.03.008>

Received 21 November 2023; Received in revised form 6 February 2024; Accepted 8 March 2024

Available online 19 March 2024

1569-1993/© 2024 European Cystic Fibrosis Society. Published by Elsevier B.V. This article is made available under the Elsevier license (<http://www.elsevier.com/open-access/userlicense/1.0/>).

## ARTICLE INFO

## Keywords:

CFTR-related disorder  
Cystic fibrosis  
CFTR  
Genetics  
Sweat test  
Nasal potential difference  
Intestinal current measurement  
Congenital absence of the vas deferens  
Pancreatitis  
Bronchiectasis  
ABPA  
Rhinosinusitis  
Sclerosing cholangitis  
Aquagenic wrinkling  
Aquagenic palmoplantar keratoderma

## ABSTRACT

This is the third paper in the series providing updated information and recommendations for people with cystic fibrosis transmembrane conductance regulator (CFTR)-related disorder (CFTR-RD). This paper covers the individual disorders, including the established conditions - congenital absence of the vas deferens (CAVD), diffuse bronchiectasis and chronic or acute recurrent pancreatitis - and also other conditions which might be considered a CFTR-RD, including allergic bronchopulmonary aspergillosis, chronic rhinosinusitis, primary sclerosing cholangitis and aquagenic wrinkling. The CFTR functional and genetic evidence in support of the condition being a CFTR-RD are discussed and guidance for reaching the diagnosis, including alternative conditions to consider and management recommendations, is provided. Gaps in our knowledge, particularly of the emerging conditions, and future areas of research, including the role of CFTR modulators, are highlighted.

The European Cystic Fibrosis Society (ECFS) and its Standards of Care Committee and Diagnostic Network Working Group have launched a project to update recommendations on the diagnosis and management of cystic fibrosis transmembrane conductance regulator (CFTR)-related disorders (CFTR-RD). Two of the four planned documents addressing the general considerations of the definition and management of CFTR-RD have been published in the *Journal of Cystic Fibrosis* [1,2]. This third paper of the series examines in more depth the characteristics of the well established disorders that are either occasionally or commonly related to CFTR dysfunction - these are congenital absence of the vas deferens (CAVD), disseminated bronchiectasis and chronic or acute recurrent pancreatitis. Conditions which might be considered a CFTR-RD, but where the mechanistic link with CFTR dysfunction is less well established, are also discussed, including allergic bronchopulmonary aspergillosis (ABPA), chronic rhinosinusitis, primary sclerosing cholangitis and aquagenic wrinkling. An important objective of this work is to not only provide guidance to CF teams, but also, importantly, to raise awareness and provide guidance to specialists outside of the CF field who are likely to encounter these conditions (e.g. experts in fertility, respiratory medicine, gastroenterology, otorhinolaryngology, dermatology, etc.).

For each disorder an overview of the clinical and pathological features is included, along with alternative aetiologies to be considered and ruled out. The key principle is that it is expected that the more common causes of each condition are excluded first by the specialist in their field, then referral to a CF service, for sweat testing and genetic analysis, should be made, so that 1) CF is excluded, and 2) CFTR-RD is confirmed (or excluded). The timing and extent of genetic analysis may vary in some situations, so further discussion is provided in each section, as is the role of further CFTR functional testing, such as nasal potential difference (PD) and intestinal current measurement (ICM). A more detailed overview of the CFTR-RD diagnostic pathway and biomarkers is provided in the first two papers of the series [1,2], although the key CFTR criteria are summarised below (Table 1).

Table 1

Key diagnostic criteria for CFTR-RD after CF has been excluded [1].

|                                                                                                                                                                                                                        |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>In vivo</i> or <i>ex vivo</i> CFTR dysfunction in the CFTR-RD range in at least 2 different CFTR functional tests (sweat test, nasal potential difference (PD) measurements, intestinal current measurements (ICM)) |
| OR                                                                                                                                                                                                                     |
| One CFTR variant and evidence of <i>in vivo</i> or <i>ex vivo</i> dysfunction in at least two functional tests                                                                                                         |
| OR                                                                                                                                                                                                                     |
| Two CFTR variants shown to reduce CFTR function, with at most one CF-causing variant                                                                                                                                   |

## 1. Congenital absence of the vas deferens

### 1.1. Veerle Vloeberghs, Dilip Nazareth, Nicholas J. Simmonds

The vas deferens (VD) is a tubular structure that connects the epididymis to the ejaculatory duct. The proximal part is palpable in the scrotum and the distal part passes through the inguinal canal and retroperitoneally into the pelvis ending in the ampulla. An absent VD can be associated with anomalies of the seminal vesicles (SV) in the form of agenesis or hypoplasia, absence of the corpus or tail of the epididymis and/or absence of the vas ampulla.

Congenital absence of the VD (CAVD) has different clinical manifestations: it can be bilateral (CBAVD) or unilateral (CUAVD), with further classification dependent on the partial or complete absence of the VD. CAVD is reported in 0.1 % of males and in up to 2 % of infertile males; it is almost universal in men with an unequivocal diagnosis of CF [3]. CUAVD is probably underestimated as conception is still possible due to normal function of the contralateral vas deferens [4]; however, a prevalence of around 1 % in males is often reported [3]. In azoospermic males, the prevalence of CBAVD is 6–12 % [5,6].

### 1.2. Diagnosis of absent vas deferens

The diagnosis of CAVD can be made (1) during systematic work-up of CF or CFTR-RD (in symptomatic patients), (2) in healthy infertile males with obstructive azoospermia, and (3) in fertile men with CUAVD discovered as an incidental finding during inguinal hernia surgery or vasectomy [7]. Incidental identification aside, the diagnosis of CAVD is based on clinical examination and semen analysis. The VD is considered absent if there is no firm, tubular structure in continuation of the epididymis that can be followed to the external inguinal canal [3]. Unfortunately, diagnosis by clinical examination is easy to miss [8]; it is often easier to be confident that the VD is present rather than absent. Only the intrascrotal part of the VD can be palpated, and sometimes a misleading structure (e.g. the fibrotic cord) can be palpated. To overcome these limitations, recent guidelines include imaging in the work-up [7–13]. By scrotal and transrectal ultrasound (TRUS), the VD, including its pelvic tract, can be evaluated, along with the SV and vas

ampulla [8,9,12,13]. Magnetic resonance imaging (MRI) can be helpful, but it is not yet established in clinical practice [14].

If CBAVD is associated with bilateral absence of the SV, semen analysis typically shows a low volume (<1 ml), low pH (<7.0) and low level of seminal plasma biochemical markers [7].


1.3. Diagnostic criteria for CFTR-RD CAVD

The most common route to diagnosis is a male presenting to a fertility specialist (andrologist/urologist) with azoospermia or oligozoospermia on semen analysis. The specialist determines the underlying pathology and anatomical abnormality, rules out other causes and considers the likelihood of CFTR dysfunction (Fig. 1). Genetic causes

other than CFTR may need consideration, such as mutations in the ADGRG2 gene [15]. In the absence of alternative aetiologies, the patient should be referred to a CF clinic (or a clinic with a specialist interest in CFTR-RD) for further evaluation to 1) exclude CF, 2) determine if CFTR-RD is the cause, and 3) to investigate for evidence of CFTR dysfunction in other organs [1].

Occasionally, CAVD will be diagnosed directly by the CF clinic, e.g. when a male is undergoing investigations for CF or CFTR-RD due to other symptoms. In this situation, semen analysis is recommended first (usually performed via fertility or urology services), followed by imaging (looking for the presence of the VD), as physical examination expertise of the VD in the CF center setting may be limited.

Individuals with CUAVD exhibit a higher frequency of CFTR variants



CF: Cystic Fibrosis, CFTR-RD: CFTR-related disease, CAVD: congenital absence vas deferens, US: ultrasound, TRUS: transrectal ultrasound, CFTR: cystic fibrosis transmembrane conductance regulator, ART: assisted reproductive technologies, ICSI: intracytoplasmic sperm injection, TESE: testicular sperm extraction, TESA: testicular sperm aspiration, FNA: fine needle aspiration, PESA: percutaneous epididymal sperm aspiration, MESA: microsurgical sperm aspiration

Fig. 1. Recommended diagnostic and management pathway for congenital absence of the vas deferens (CAVD).

than healthy men. At least one *CFTR* variant is found in 46 % of CUAVD males, with *5T*, *R117H* and *F508del* being the most common variants [16]. While *5T* is a relatively frequent variant [17], in combination with (TG)12 or (TG)13 repeats in *CFTR* intron 9, it may increase CAVD severity (e.g. increase the risk of partial to complete obstruction/absence of the VD) [18]. These individuals should be worked up in a similar fashion to those with CBAVD.

#### 1.4. Alternative causes of CAVD

Two different mechanisms have been suggested to be responsible for CAVD. The first one is atrophy-based where dysfunction of the seminal duct epithelium disrupts the intraluminal fluid homeostasis, leading to thick secretions and involution of the VD. Alternatively, CAVD might be the result of VD agenesis during the embryonic development of the Wolfian duct [7].

Renal abnormalities (25 %), often in the form of ipsilateral renal agenesis, may be detected during CAVD assessment [19]. CUAVD is more frequently associated with unilateral renal absence than CBAVD - 39 % and 19 %, respectively [11]. Men carrying *CFTR* mutations are unlikely to have renal agenesis [20].

#### 1.5. Recommendation for follow-up and treatment of CFTR-RD CAVD

Isolated CFTR-RD CAVD is not usually considered a progressive (or reversible) condition, but individuals should be made aware that over time other CFTR-related manifestations might develop. As it is unreasonable to expect an andrologist to follow-up a patient with CFTR-RD long-term, annual follow-up in a CF clinic or a clinic specific for CFTR-RD is recommended, as the lifetime risk of developing other CFTR-related complications is currently unknown. Alternatively, if follow-up is not wanted/feasible, the person with CAVD and their family physician should be made aware of the clinical features to look out for which would trigger a referral back to the CF center for further evaluation [1].

## 2. Disseminated bronchiectasis

2.1. Peter Middleton, James Chalmers, Damian Downey, Jerry Nick, Kris De Boeck

Bronchiectasis is characterised by abnormal dilatation of the large and medium airways, often also involving the small airways [21]. Affected airways become inflamed, thick walled and irregularly dilated, resulting in airflow obstruction and impaired clearance of secretions. Bronchiectasis can present as a focal process in one airway due to partial or complete obstruction of the airway (e.g. tumour, inhaled foreign body) or following severe pneumonia. More commonly, bronchiectasis is a generalised disease involving many airways in different lobes, causing recurrent lower respiratory tract infections, chronic cough and sputum production.

#### 2.2. Diagnosis of bronchiectasis

Bronchiectasis is defined as an increase in the size of an airway relative to the accompanying pulmonary artery, generally measured on High Resolution Computerized Tomography (HRCT). Since airway size increases with increasing lung volume, the level of inspiration should be standardized during HRCT. The Bronchus to Artery Ratio (BAR) increases with age, with an upper limit of normal in young children of around 0.76, increasing up to 1.0 in adults [22,23].

The possibility of bronchiectasis should be considered in people with chronic “wet” cough with or without expectoration. A new diagnosis of bronchiectasis should prompt a thorough investigation in a specialist service to find the underlying cause [24,25].

#### 2.3. Diagnostic criteria for CFTR-RD bronchiectasis

The diagnosis of CFTR-RD should be considered for people with bronchiectasis, especially in the absence of other causes (see below), when there is evidence of CFTR protein dysfunction but the diagnostic criteria for CF are not fulfilled, as outlined in the first paper of this CFTR-RD series [1]. In settings where CFTR functional tests, such as intestinal current measurement (ICM) or nasal potential difference (PD), are not available, it may be helpful to first search for *CFTR* variants by genetic testing before referring the patient [1]. This is especially important in people with bronchiectasis and intermediate sweat test results. Historical sweat tests reported narratively as ‘normal’ or ‘borderline’ should be repeated.

An association between *CFTR* variants and bronchiectasis has long been reported. Uncontrolled, observational studies have reported at least one *CFTR* variant in up to 50 % of people with bronchiectasis, with a high incidence of the *5T* splicing variant of the poly T tract [26–28]. In one study, two sequence variants were found in up to 20 % of cases, though segregation analysis was not always performed to ensure the variants were inherited *in trans*. Other studies, however, have not shown an association [29]. A population-based investigation examined more than 100,000 carriers of the *F508del* variant, showing that carriers had an odds ratio of 1.31 (95 % confidence interval (CI) 1.16–1.48) for chronic bronchitis and a hazard ratio of 1.88 (95 % CI 1.03–3.45) for bronchiectasis [30].

The involvement of CFTR dysfunction in bronchiectasis associated with other systemic diseases raises further considerations. An extensive genetic, clinical and physiological examination of 26 people with rheumatoid arthritis and bronchiectasis reported that seven had sequence variants in *CFTR*, including four with *F508del* [31]. Of the remaining 19 individuals, seven had sweat chloride values  $\geq 60$  mmol/L. Only two of 25 sweat tests demonstrated chloride values in the normal range ( $<30$  mmol/L). Importantly, 20 of the 26 individuals had childhood onset bronchiectasis often predating the onset of rheumatoid arthritis by decades, highlighting the difficulty of assigning causation at an individual point in time [31]. Similarly for people with a diagnosis of bronchiectasis due to ABPA, an increased incidence of *CFTR* variants has been reported (discussed later in this paper) [32]. These data indicate that in certain circumstances it may be appropriate to revisit an assigned diagnosis (such as rheumatoid arthritis-associated bronchiectasis or ABPA) to determine if these individuals now fulfil the updated diagnostic criteria for CFTR-RD [1].

#### 2.4. Alternative causes of disseminated bronchiectasis

In addition to ABPA and rheumatoid arthritis, disseminated bronchiectasis can be associated with other underlying conditions, including a variety of immune deficiencies, primary ciliary dyskinesia (PCD), autoimmune diseases, inflammatory bowel disease and other connective tissue disorders. Significant infections may also result in bronchiectasis, termed “post-infectious,” with the remaining cases termed “idiopathic,” although there is emerging evidence that *CFTR* variants (without CF or CFTR-RD) may alter this risk [33]. Despite an exhaustive evaluation, a specific aetiology may not be identified in up to half of cases [24,25].

A detailed medical history should focus on the age of onset of symptoms, in particular neonatal respiratory distress and early childhood respiratory infections, weight loss, hearing problems, infertility, presence of nasal polyps or chronic ear infections and gastrointestinal symptoms, including gastro-oesophageal reflux, and a family history [24]. Routine testing should include differential blood count, serum immunoglobulins and testing for ABPA (see below for more details on ABPA testing) [24]. Depending on history, findings on physical examination and HRCT, other tests may be indicated.

The distribution of bronchiectasis on HRCT can point towards a specific aetiology, with upper lobe predominant disease found in CF and lower and middle lobe predominant disease in PCD and immune

deficiencies, in some but not all studies [34–36]. Other aspects may point towards a diagnosis of CF, including airway infections with *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Burkholderia* spp. and non-tuberculous mycobacteria (e.g. *Mycobacterium abscessus* and *M. avium*). Extrapulmonary manifestations may also increase the suspicion for CF, including male infertility, pancreatitis and/or exocrine pancreatic insufficiency, bowel disease, chronic sinusitis and/or nasal polypoidosis and skin changes (such as a salt crust following exercise or aquagenic wrinkling).

The diagnostic tests for CF and CFTR-RD have already been outlined above, but the assessment for other CFTR-related organ pathology may also be indicated (e.g. pancreatic function and semen analysis). More specialised measures of CFTR function such as ICM and nasal PD is recommended in people with intermediate results on sweat testing and CFTR analysis. Adults presenting with bronchiectasis and characteristic involvement in at least one other organ may have CF and severe lung disease may already be present at the time of diagnosis [37,38].

The “PICADAR” (PrIMary Ciliary Dyskinesia Rule) scoring system for PCD may assist in the assessment of people with chronic sputum production in childhood [39]. A combination of nasal nitric oxide testing, ciliary structure/function testing (by electron microscopy and high speed video microscopy) and genetics should be considered to investigate for PCD [40].

Testing of autoantibodies to screen for connective tissue or autoimmune disease, is indicated in the presence of suggestive findings towards these pathologies on history and physical examination. The diagnosis of post-infectious bronchiectasis is typically reserved for those with severe lung infections earlier in life and is rare in individuals with access to standard medical care.

## 2.5. Recommendations for follow-up and treatment of CFTR-RD bronchiectasis

The European Respiratory Society (ERS) and British Thoracic Society (BTS) published guidelines for the treatment and follow up of adults with non-CF bronchiectasis with separate standards of care for CF [24, 25]. The first paper in this series provides an overview of the general recommendations for the follow-up of people with CFTR-RD but, in the absence of specific guidelines for people with CFTR-RD bronchiectasis, the ERS/BTS guidelines can be considered, recognising that many recommendations lack a strong evidence-base. Individualised treatment based on clinical acumen should achieve symptom control, improve quality of life and prevent pulmonary exacerbations. This is likely best achieved in specialist centres, ideally attached to CF centres, including clinics specific for CFTR-RD.

The recent BTS guidelines suggest a follow-up frequency appropriate to the patients’ disease severity with at least an annual specialist review during clinical stability for assessment of symptoms, exacerbation frequency and presence of comorbidities, as well as for measurements of body mass index, dyspnoea score, pulse oximetry, sputum bacteriology culture and lung function [25]. Review by a specialist respiratory physiotherapist is recommended and airway clearance techniques should be initiated for chronic productive cough or difficulty with sputum expectoration. Subjects with impaired exercise capacity should participate in a pulmonary rehabilitation programme and take regular exercise.

When standard airway clearance techniques do not control symptoms, mucoactive treatment such as hypertonic saline should be considered. Although recombinant human DNase is contraindicated in non-CF bronchiectasis overall, given the beneficial effects in people with CF, use may be evaluated on a case-by-case basis in those with CFTR-RD. Long acting bronchodilators and inhaled corticosteroids, are useful in subjects with concomitant asthma or COPD but are not generally recommended for use in people with bronchiectasis [24].

Antibiotic eradication treatment should be offered to patients with a new sputum isolation of *P. aeruginosa*, but is not generally recommended

for other pathogens [24]. Long-term antibiotic treatment is indicated in patients who experience three or more exacerbations per year, since both macrolides and inhaled antibiotics have been proven effective to reduce exacerbation frequency in randomized controlled trials of people with bronchiectasis [25].

Pulmonary exacerbations should be treated with oral/inhaled/intravenous antibiotics as per the CF and non-CF treatment guidelines [24,25,41,42]. Currently CFTR modulators are not approved outside of the CF diagnosis, although as CFTR dysfunction is present in CFTR-RD bronchiectasis, it seems highly plausible that benefit will be derived for many, although further research is required.

In summary, individuals with CFTR-RD bronchiectasis should be counselled appropriately about their condition, explaining that our understanding of CFTR and its relationship to bronchiectasis has evolved, with discussion about why the distinction between CF and CFTR-RD has been made (see paper 1 of this series for advantages/disadvantages) [1]. Until evidence specific for subjects with bronchiectasis due to CFTR-RD emerges, individualised treatment should be based on the guidelines for both CF and non-CF bronchiectasis, taking into account both disease severity and other factors pertinent to that individual.

## 3. Chronic or acute recurrent pancreatitis

3.1. Frank Bodewes, Michael Wilschanski, Isabelle Scheers, Chee Y. (Keith) Ooi, Isabelle Sermet-Gaudelus

Acute pancreatitis (AP) is characterised by acute inflammation of the pancreas and is associated with typical symptoms suggestive of pancreatic origin (e.g. severe abdominal pain, nausea and vomiting) and elevated serum amylase or lipase  $\geq 3$  times the upper limit of normal and/or imaging findings (e.g. ultrasound) compatible with acute pancreatitis. Acute recurrent pancreatitis (ARP) is characterized by recurrent episodes of inflammation of the pancreas. The diagnosis requires at least two episodes of acute pancreatitis with complete resolution of pain and normalization of amylase and lipase between episodes. Chronic pancreatitis (CP) is a progressive and irreversible inflammation of the pancreas defined by at least one of the following three criteria: 1) abdominal pain of pancreatic origin with imaging findings of chronic pancreatitis (e.g., calcification); 2) evidence of exocrine pancreatic insufficiency with imaging findings of CP; and, 3) endocrine pancreatic insufficiency with imaging findings of CP. All types of pancreatitis can be caused by known or unknown underlying conditions (see below). CP can lead to permanent loss of function of the pancreas and requires lifelong follow up and management.

CFTR plays a crucial role in the physiological functioning of the pancreas. Defects in CFTR lead to impaired bicarbonate secretion, disrupting the normal balance of pancreatic secretions and contributing to the development of pancreatitis and other pancreatic diseases. Diagnostic criteria for CFTR-RD pancreatitis rely on (1) excluding other aetiologies and (2) confirming the criteria for CFTR-RD [1].

## 3.2. Diagnosis of pancreatitis

In addition to measuring serum amylase and lipase levels, the initial workup for ARP typically includes full liver enzymes, calcium, fasting triglycerides, coeliac antibodies (e.g. anti-TTG [tissue transglutaminase]) and IgG4, pancreatic imaging (e.g. Magnetic Resonance Cholangiopancreatography, MRCP), sweat chloride analysis and genetic testing [43,44]. Apart from CFTR, there are other genes associated with an increased risk of developing pancreatitis, including variants in *PRSS1*, *SPINK1*, *CTRC*, *TRPV6* and *CPA1*, amongst others [45–47]; these risk factors may be present even among those who carry 1–2 CFTR variants. Additional testing to consider include serum amino acids and urine organic acids (for the inherited conditions methylmalonic academia and propionic academia) and stool for parasites/fungi as these

are all rare but important causes of pancreatitis

Anatomical anomalies, specifically pancreas divisum, are associated with ARP in 5–20 % of cases. In ARP and CP, MRCP is the preferred imaging technique for the assessment of the pancreas. Endoscopic cholangiopancreatography (ERCP) is usually reserved for therapeutic purposes.

3.3. Alternative causes of pancreatitis

The causes of AP, ARP and CP are diverse – internationally recognised ‘checklists’ of the causes include TIGAR-O (Toxic-metabolic, Idiopathic, Genetic, Autoimmune, Recurrent and severe acute pancreatitis and Obstructive) checklist for adults, and INSPPIRE (International Study Group of Pediatric Pancreatitis: In Search for a Cure) checklist for children [43,44,48]. The evaluation should thus include a history of possible past attacks, medications taken, exposure to toxins, a family history of pancreatitis, diabetes, alcohol, and gallbladder disease. Primary aetiologies of pancreatitis should be carefully investigated, including toxic or metabolic causes (e.g. alcohol, tobacco, hypercalcaemia, hyperlipidemia, medications), autoimmune factors, and obstructive causes [43,49]. Some of these may only be risk factors, which increase the likelihood of developing pancreatitis but do not directly cause it. Risk factors for pancreatitis include genetics, lifestyle, and environmental.

3.4. Diagnostic criteria for CFTR-RD pancreatitis

As outlined above, pancreatitis is widely recognized as a multifactorial disease, resulting from the combined impact of multiple insults, including genetic variants and anatomical factors, like a pancreas divisum [43,48]. All of these need to be considered. Patients at risk of pancreatitis usually have normal exocrine pancreatic function, and one of the CFTR variants in these pancreatic sufficient patients is usually associated with residual function.

To determine if a patient has AP, ARP, or CP associated with CFTR protein dysfunction, it is important to assess for CFTR variants and protein function [1,2]. Next-Generation Sequencing for pancreatitis gene variants should be performed in all patients with otherwise unexplained ARP or CP, including the CFTR polyT variants. If CFTR variants are identified, their relevance should be assessed by evaluating CFTR function primarily via a sweat test, but other functional tests such as nasal PD and ICM may be required. It is important to note that a normal sweat chloride (<30 mmol/L) does not rule out the diagnosis of CF in all cases (even though it is considered standard care for ARP and CP). The order of genetic testing and CFTR function testing, particularly sweat

testing, can differ depending on the individual situation and circumstances (Table 2).

CFTR dysfunction can affect chloride and bicarbonate transport. Several reports have suggested that specific CFTR variants (e.g. R74Q, R75Q, R170H, L967S, R1162L, L997F, D1152H, S1235R, D1270N and R117H) can lead to impaired bicarbonate secretion and result in various disease states, including pancreatitis [50,51]. Some CFTR variants that alter bicarbonate but not chloride transport have been associated with a normal sweat test [50], which reinforces the need to always perform genetic testing in this clinical context. Indeed, the current definition of CFTR-RD only includes CFTR functional tests (sweat test, nasal PD, ICM) related to chloride transport. Therefore, these functional tests may not detect a bicarbonate transport defect, and genetic testing is mandatory in this clinical situation to differentiate pancreatitis in CFTR-RD from CF and CFTR carriers (Table 2).

3.5. Recommendations for follow-up and treatment of CFTR-RD pancreatitis

The management and treatment of pancreatitis in general is applicable to this population and is beyond the scope of this article. The issues specific to CFTR function are summarised in Table 2. The impact of CFTR modulators on pancreatic function and related complications have been reported [52,53]. In people with CF, ivacaftor has been shown to improve symptoms and reduce inflammation in CF pancreatitis [54], and CFTR modulators have been linked to a decrease in hospitalizations due to acute pancreatitis [55]. Conversely, it has also been suggested that CFTR modulators may increase the incidence of pancreatitis in exocrine pancreatic insufficiency [56]. Currently, classifying pancreatitis as CFTR-RD does not impact standard disease treatment, as CFTR modulators are not approved beyond the strict diagnosis of CF. Since the CFTR-RD classification for pancreatitis provides a mechanistic link between CFTR mutations and CFTR protein-related chloride (or bicarbonate) transport in individual patients, CFTR modulators might in the future be considered as potential treatment option for CFTR-RD-associated pancreatitis. Further research is needed to establish the effects of CFTR modulators, determine eligibility, optimal dosage and treatment duration, across the full spectrum of CFTR function, so that an accurate assessment of risk/benefit can be determined.

**Table 2**  
Pancreatitis and CFTR: a summary of the different clinical scenarios and recommendations.

| Acute recurrent (>1 episode) and chronic pancreatitis |                                                                                                                                                                                                                                                                         |                                                                                                                                                                                      |                                                                                                                                                                                                      |                                                                                       |
|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------|
| Diagnosis                                             | Cystic fibrosis with pancreatitis                                                                                                                                                                                                                                       | Pancreatitis in CFTR-RD                                                                                                                                                              | Pancreatitis in CFTR carriers                                                                                                                                                                        | Pancreatitis with no CFTR variants                                                    |
| Key diagnostic tests                                  | Exclude common causes of pancreatitis<br>Newborn screening<br>CFTR genetic testing*<br>Sweat test                                                                                                                                                                       | Exclude common causes of pancreatitis<br>CFTR genetic testing*<br>Sweat test<br>Nasal PD<br>ICM                                                                                      | Exclude common causes of pancreatitis<br>CFTR genetic testing*<br>Sweat test<br>Consider Nasal PD<br>Consider ICM                                                                                    | Exclude common causes of pancreatitis<br>CFTR genetic testing* (no variants detected) |
| Key CFTR-related management principles                | Patient diagnosed with CF developing pancreatitis or CF patients with pancreatitis as the primary presentation should be treated as a CF patient for ongoing care and management.<br>Refer to CF specialist center.<br>Potentially eligible for CFTR modulator therapy. | Diagnose CFTR-RD according to the ECFS CFTR-RD diagnostic guidelines.<br>Refer to CF specialist center.<br>Theoretically might be eligible for CFTR modulator therapy in the future. | When a single CFTR mutation is detected, it is necessary to exclude CFTR-RD (and CF).<br>Refer to CF specialist center.<br>Theoretically might be eligible for CFTR modulator therapy in the future. | CFTR does not play a role in either causing or increasing the risk of pancreatitis.   |

When the diagnosis is unclear, consider testing for a panel of genes associated with pancreatitis (see text).  
Abbreviations: CFTR=cystic fibrosis transmembrane conductance regulator; PD=potential difference; ICM=intestinal current measurement; ECFS=European CF Society. \*Extended CFTR analysis should be performed if first-line targeted assessment is inconclusive.

#### 4. Chronic Rhinosinusitis

4.1. Jochen G. Mainz, Assen Koitschev, Daniel Peckham, Donald Vandevanter, Kevin W Southern

Chronic rhinosinusitis (CRS) is a common condition with a number of underlying causes affecting children and adults. It is a characteristic feature of CF, especially in adults, and is a significant cause of morbidity in this population [57]. Nasal polyposis, occasionally extensive, is also a feature of CF. In contrast, aside from medication toxicity (e.g. aminoglycosides, azithromycin) of the vestibulocochlear system secondary to treatment of CF airway bacterial pathogens, the ear does not appear to be affected by CFTR dysfunction. An increased prevalence of chronic rhinosinusitis has also been reported in carriers of *CFTR* variants [58, 59].

Rhinosinusitis is inflammation of the mucosa of the nasal and paranasal cavities characterised by two or more symptoms, one of which should be either:

- nasal blockage / obstruction / congestion
- or
- nasal discharge (anterior and/or posterior nasal drip)

Other symptoms can include facial pain/pressure and/or reduction or loss of smell as a second diagnostic symptom according to the European Position Paper on Rhinosinusitis and Nasal Polyps 2020 [60].

For a CRS diagnosis, the above mentioned symptoms should persist for more than 12 weeks. The extent of CRS can be confirmed either by endoscopic examination, and/or CT or MRI imaging of the sinuses [61]. Typical endoscopic findings include nasal polyps and/or mucopurulent discharge and/or oedema/mucosal obstruction, primarily from the middle meatus. Pathological changes observed on CT/MRI are most frequent within the ostiomeatal complex and maxillary sinus.

#### 4.2. Can CRS be considered a CFTR-RD?

Given that CRS is a hallmark of CF, that virtually all adults with CF have some sinonasal abnormalities by imaging, and that the presence of nasal polyps in preschool children are highly predictive of CF, it seems reasonable to assume that the upper airway is sensitive to CFTR dysfunction and that CRS is a candidate condition for CFTR-RD [61–65]. The resolution of CRS symptoms among those treated with CFTR modulators supports the rationale that CFTR dysfunction is causative [65–67]. A study of family members related to people with CF found an increased prevalence of CRS, supporting the assertion that mildly reduced CFTR function may be associated with CRS, even among obligate CF carriers [68]. A case-control study comparing people with CRS to age matched controls found an increased prevalence of CF-causing *CFTR* variants, again supporting the hypothesis that for some people with CRS, the underlying cause may be CFTR dysfunction and that the condition represents a CFTR-RD [68].

#### 4.3. CFTR-RD CRS diagnostic criteria and recommendations for follow-up and treatment

To fulfil the diagnostic criteria for CFTR-RD, alternative causes of CRS and a diagnosis of CF should both be excluded [1].

#### 4.4. Alternative causes of CRS

CRS is frequently associated with inhaled allergens [60]. Allergic secretions are usually serous and often combined with endoscopic findings of pale swollen mucosa and associated with other allergic manifestations e.g. conjunctivitis. The histological appearances of allergy-associated nasal polyps is a predominant eosinophilic inflammatory infiltrate, which typically responds well to topical corticosteroid

therapy.

CRS is a hallmark of PCD, which is associated with impaired mucociliary clearance due to ciliary dysfunction. The pattern of PCD airway bacterial colonization is similar to that of CF or CFTR-RD, although *P. aeruginosa* is detected less frequently. PCD is also associated with situs inversus and troublesome chronic otitis media, which can lead to difficulty in hearing and speech delay [69].

#### 4.5. Excluding a diagnosis of CF

CF should be excluded in individuals with severe CRS, with or without nasal polyps, refractory to anti-allergic therapies, especially when associated with purulent secretions and isolation of CF-typical bacterial pathogens from the airways, such as *P. aeruginosa*. As CRS is relatively common, sweat testing first is recommended. Only if refractory disease persists, despite repeated sinonasal surgery and/or the sweat test is not normal (<30 mmol/L), should more extensive testing (including genetic analysis) be performed, following ECFS CFTR-RD diagnostic guidelines [1]. Furthermore, in pre-schoolers, the presence of nasal polyps (not hyperplastic adenoids, which may be erroneously termed ‘polyps’) should prompt CFTR investigations. In some cases, more extensive electrophysiological testing may be appropriate (e.g. ICM or nasal PD, although it should be noted that significant nasal inflammation can affect the quality of nasal PD measurements) [70]. Genetic testing may identify variants associated with CFTR-RD or carriers of *CFTR* variants. A negative genetic test does not exclude CFTR-RD, but makes it less likely, especially if other potential CRS causes are identified [1,2]. With some evidence of CFTR dysfunction and/or identification of *CFTR* variants, in the absence of another cause of CRS, a diagnosis of CFTR-RD seems appropriate.

#### 4.6. Recommendations for follow-up and treatment of CFTR-RD CRS

A history of nasal blockage, snoring and mouth-breathing is indicative of CRS, together with purulent secretions visible by rhinoscopy and often also by oral inspection, e.g. on the dorsal pharyngeal wall from postnasal drip. If questioned or assessed by methods such as “Sniffin sticks,” impaired smelling, reduced taste and appetite, may be detected [71]. Monitoring of progress should include direct visualisation of the nasal cavity with nasendoscopy and imaging. In absence of a nasendoscope, the examination can be performed by an otoscope usually available to physicians in routine practice.

In patients with CRS refractory to conservative therapies, sinonasal imaging should be performed, ideally by MRI to monitor progress and need for surgical intervention. Sinus radiographs are of low diagnostic value and are associated with radiation exposure (which is not associated with MRI), so should be avoided. Low-dose CT scans can be performed on an individual basis, especially before scheduled surgery. Diagnostic nasal lavage [72], such as by nasal passage rinsing with 10mls of isotonic saline or nasal swabs [73], facilitates detection of CF-typical bacteria (e.g. *P. aeruginosa*) in the sinonasal space, which has been identified as a reservoir and a site of first colonization of the “united airway” system. However *P. aeruginosa* may also be isolated in other diseases, such as PCD, with similar clones being identified in both the lungs and paranasal sinuses [74].

CRS is actively treated in CF with topical nasal therapies and surgical intervention. The extent of treatment depends on the impact of the condition on the quality of life of people with CF (and also likely CFTR-RD). The management of CF-related CRS depends on a good working partnership between CF and ENT teams [75]. A similar active ethos is appropriate for patients with CFTR-RD. A systematic review of therapies for CF-related CRS identified clear evidence to support the use of nasal douches with saline, endoscopic sinus surgery and CFTR modulators [76]. Evidence for topical corticosteroids and antibiotics was less robust but these interventions should be considered [76,77]. Importantly, inhalation with conventional aerosols and even nasal lavages do not

sufficiently reach the paranasal sinuses, but this can be achieved with pulsating aerosols [78]. This is particularly important if attempting to eradicate pathogens from the paranasal sinuses. Small studies have demonstrated some effectiveness of mucolytics and antibiotics, but there is a lack of large trials in this field [79–81].

Conservative therapies for upper airway blockage by polyps and/or swollen mucosa include topical nasal steroids as first-line long term therapy, even if sufficiently powered studies have not yet been performed in CF [82]. For relieving the airways from mucopurulent secretions and crusts, therapeutic nasal lavages are performed with 250mls of isotonic or slightly hypertonic saline.

Surgery is required if conservative therapeutic measures are unsuccessful. However, relapses are frequent, as the underlying defect remains. CFTR modulators for people with CF and significant CRS have been shown to reduce sinonasal inflammation, as well as improvements in imaging and endoscopy findings [65,66,83,84]. On rare occasions, CFTR modulators are also prescribed for non-pulmonary indications, including sinus disease, in people with CF after lung transplantation [85]. Further studies are needed to characterise the relative risk/benefit of such treatment in this group. All of the aforementioned treatment strategies should also be relevant to CFTR-RD, although specific evidence is lacking. Importantly, there is currently no evidence to support the use of modulators in CFTR-RD CRS, but as these agents become generic and more accessible, a role may emerge.

## 5. Allergic bronchopulmonary aspergillosis

### 5.1. Carsten Schwarz, Patience Eschenhagen, Michael Waller

Allergic bronchopulmonary aspergillosis (ABPA) is an immunological disease triggered by inhalation of *Aspergillus* spp. spores in susceptible and genetically predisposed individuals, such as people with asthma or CF [86,87]. It is characterised by a T-helper-2 cellular response and humoral response resulting in the production of polyclonal IgE as well as *Aspergillus*-specific IgE, IgG and IgA antibodies. The clinical presentation is usually an acute or subacute clinical deterioration with cough, wheeze, exercise intolerance, exercise-induced asthma, decline in pulmonary function, and increased sputum. The CF community relies mainly on the consensus criteria issued by the Cystic Fibrosis Foundation in 2003 (Table 3) [88].

### 5.2. Can ABPA be considered a CFTR-RD?

The actual prevalence of ABPA in CF is difficult to establish because of the substantial overlap with classical CF respiratory disease. Retrospective data from epidemiological studies report a range of 2–14 % [89, 90]. Pathophysiological mechanistic links between ABPA and CFTR deficiency are described in the literature. In one systematic review and meta-analysis a possible pathogenetic link between CFTR variants and ABPA was shown as the odds of encountering CFTR variants was higher in ABPA compared with the control group (odds ratio (OR) 10.39; 95 % CI, 4.35–24.79) or the asthma population (OR 5.53; 1.62–18.82) [32]. In an ABPA patient cohort [91], *F508del* was identified more often than in the general population and another study reported a significantly higher frequency of CFTR heterozygotes among ABPA patients (28.5 %) compared to control asthmatic patients (4.6 %;  $p = 0.01$ ) and individuals seeking genetic counseling ( $p < 0.001$ ) [92]. A further study found the

frequency of *F508del* to be higher in patients with ABPA than in 53 Caucasian patients with chronic bronchitis ( $p < 0.0003$ ) and the general population ( $p < 0.003$ ) [93].

A link between CFTR and ABPA is also supported by evidence from animal models. The *Aspergillus fumigatus* extract sensitisation of CFTR-Null mice has been reported to affect eicosanoid pathway gene expression with distinct regulation of *PLA2G4C*, *PLA2G2D* and *ALOX15* genes, thus impacting ABPA and CF [94]. CFTR-Null mice also showed a significantly elevated response in IgE to *Aspergillus* sensitisation as well as a shift from IL-5 to IL-4 responses compared to controls [95]. Furthermore, gene therapy with rAAV5Delta-264CFTR attenuated the hyper-IgE response in a reproducible CF mouse model of ABPA, with systemic effects also evident in the cytokine response of stimulated splenocytes [96]. These data substantiate a correlation between CFTR dysfunction and ABPA and suggest an aetiological role of CFTR in a subset of ABPA patients. In these individuals ABPA may be considered a CFTR-RD.

### 5.3. CFTR-RD ABPA diagnostic criteria and recommendations for follow-up and treatment

Before reaching a diagnosis of CFTR-RD, alternative conditions associated with ABPA, including CF itself, should be excluded. Other associated conditions include asthma, COPD and non-CF bronchiectasis [2,97–101]. After excluding these conditions, evidence of CFTR dysfunction should be sought by sweat testing and genetic analysis (followed by nasal PD and/or ICM, if indicated), as per the CFTR-RD diagnostic guidelines [1,2]. As the exclusion of CF and the interpretation of the diagnostic tests may prove challenging, the diagnosis of CFTR-RD ABPA should be made in a CF center [1]. As for the other CFTR-RD conditions, the ideal follow-up should take place in a CF clinic or specialist CFTR-RD clinic, where the possible involvement of other organs and evolution of the clinical condition are appropriately monitored [1]. As treatment of ABPA can be challenging, particularly when CFTR impairment is present, close collaboration with specialists in fungal disease is recommended.

## 6. Primary sclerosing cholangitis

### 6.1. Michael Wilschanski, Frank Bodewes, Dominique Debray, Isabelle Sermet-Gaudelus

Primary sclerosing cholangitis (PSC) is a progressive inflammatory and scarring disease of the large bile ducts that may lead to ductal obstruction, cholestasis, cholangitis, biliary fibrosis, and cirrhosis. The diagnosis is based on radiographic features. The cholangiographic features on MRCP or ERCP are specific and include the characteristic "beading" appearance of dilatation and stricturing involving the intrahepatic and/or extrahepatic biliary tree. PSC is histologically characterized by progressive periductal fibrosis with luminal stenosis or obliteration, along with the formation of a fibrous core, and dilatation (cholangiectasis). PSC is a rare disease, with a frequency of 0.6 per 100,000 patients/year and a bimodal peak at 15 and 35 years of age. It is the most common hepatobiliary disease associated with inflammatory bowel disease, particularly ulcerative colitis. Many patients are asymptomatic, and the diagnosis is suggested by routine liver tests. 10–15 % of adult patients present with anorexia, weight loss, pruritus,

**Table 3**

ABPA definition in CF [88]. These criteria are also likely applicable to CFTR-RD.

- a (sub-)acute clinical deterioration not attributable to another aetiology
- total serum IgE >1000 IU/mL in steroid-free patients
- positive immediate cutaneous hypersensitivity to *Aspergillus* antigens (Ags) or elevated *A.fumigatus*-specific IgE levels
- precipitating or IgG antibodies against *A.fumigatus* in serum
- and new radiological abnormalities, which have not cleared with physiotherapy and antibiotic treatment.

jaundice, and right upper abdominal quadrant pain.

6.2. Can PSC be considered a CFTR-RD?

The similar cholangiographic characteristics of CF and PSC have prompted investigations on the role of CFTR dysfunction in the development of PSC [102,103]. Genetic analysis of *CFTR* variants in children and adults with PSC has yielded controversial results. An increased frequency of *CFTR* pathogenic variant heterozygosity has been reported, but this was not significantly different to the rate observed in the control adult population in other studies [104–109]. Functional analysis showed in a minority of patients a decrease of CFTR function to the level of 30 to 65 % wild type [106,107,109]. Of the approximately 100 patients studied so far, a subgroup of 20 would be considered to have CFTR-RD on the basis of current diagnostic criteria [106,107,109]. These observations raise the question of whether a decrease in CFTR function can contribute to the development of PSC, and whether CFTR modulators would be of potential benefit. Larger studies involving patients with PSC tested for CFTR dysfunction are warranted to investigate this hypothesis.

6.3. Diagnostic criteria and recommendations for follow-up and treatment

Other causes of sclerosing cholangitis should be excluded before investigations for CFTR-RD are considered (Table 4). PSC is rare in adults and children, and its aetiology remains mostly unknown. Particularly in children, there is an association of PSC with autoimmune sclerosing cholangitis, characterized by the presence of autoantibodies (ANA, SMA, LKM1) and histological features of sclerosing cholangitis with other autoimmune features. Other secondary causes of sclerosing cholangitis are listed in Table 4.

Based on the available evidence, if no other cause of sclerosing cholangitis is identified, *CFTR* genetic testing is recommended. This should be followed by sweat testing and further electrophysiological testing (nasal PD and/or ICM), if indicated [1]. As per updated

**Table 4**  
Causes of sclerosing cholangitis to be considered before investigations for CFTR-RD are initiated.

| Pediatric age group [110]       | Adult age group [111]                                                                 |
|---------------------------------|---------------------------------------------------------------------------------------|
| Neonatal sclerosing cholangitis | Cryptosporidiosis, microsporidiosis                                                   |
| PFIC 3 / MDR 3 deficiency       | Ruptured hydatid (parasite)                                                           |
| Ischemic injury (ITBL)          | Recurrent pyogenic cholangitis/Oriental cholangitis                                   |
| Bacterial cholangitis           | CMV infection                                                                         |
| Portal biliopathy               | Primary immunodeficiency syndromes                                                    |
| Surgical complications          | Secondary immunodeficiency (AIDS)                                                     |
| Cystic fibrosis                 | Critical illness disease                                                              |
| AIDS-related cholangiopathy     | Hepatic allograft                                                                     |
| Retroperitoneal fibrosis        | Hepatic artery thrombosis/stenosis/ ligation                                          |
| Langerhans cell histiocytosis   | Chemotherapy                                                                          |
| Primary immunodeficiency        | Ig4 associated SC                                                                     |
| Secondary immunodeficiency      | Eosinophilic cholangitis                                                              |
| IgG4-associated cholangitis     | Mast-cell cholangiopathy                                                              |
| Eosinophilic Cholangitis        | Choledochal varices (cavernous formation)                                             |
| Sickle cell disease             | Choledocholithiasis                                                                   |
| Histiocytosis X                 | Iatrogenic stricture                                                                  |
| CMV infection                   | Malignant infiltration/Cholangiocarcinoma                                             |
| Cryptosporidium infection       | Langerhans histiocytosis                                                              |
|                                 | Caustic or alcohol instillation into bile duct, eg, treatment of hepatic hydatid cyst |

Abbreviations: CMV=cytomegalovirus; ITBL=ischemic type biliary lesions; AIDS=acquired immunodeficiency syndrome.

guidelines, if CFTR-RD is confirmed, then follow-up in a CF or specialist CFTR-RD centre is advised, in collaboration with a specialist gastroenterology team [1]. There is no known treatment for PSC. Ursodeoxycholic acid may improve liver tests but has no impact on outcome [110]. Portal hypertension may develop with progressive disease. PSC is associated with hepatobiliary malignancies, particularly cholangiocarcinoma. Liver transplantation is at present the only treatment.

7. Aquagenic wrinkling of the palms

7.1. Elke De Wachter, Emmanuelle Girodon, Caroline Raynal, Geoffroy Hickman, Emmanuelle Bourrat

Aquagenic wrinkling of the palms (AWP), also named aquagenic palmoplantar (pseudo) keratoderma (APPK), is a rare skin disease characterized by transient wrinkling, oedema and whitish papules mostly on the palms, with pruritus, burning and pain appearing within five minutes of water immersion [112,113]. It resolves rapidly after drying. The diagnosis is made by immersion of both hands in tap water for five minutes followed by clinical examination of the palms [112,114, 115]. No standardized test is available and differences occur in the literature on water temperature (range: 15–40 °C), immersion time (5–15 min) and scoring systems [114–118]. Histopathological features include thickness of the stratum corneum, however, a biopsy is not needed for the diagnosis [119].

7.2. Can AWP be considered a CFTR-RD?

AWP has been reported in up to 84 % of patients with CF [115,120], as well as 8 % of obligate CF carriers [116]. The improvement of AWP in response to CFTR modulator therapy in CF patients underlines the relationship between dysfunctional CFTR and AWP [121,122]. Only one large study, involving 112 French patients presenting with isolated AWP, has assessed the frequency of pathogenic *CFTR* variants [123]. A significant excess of *CFTR* variant carriers (33 %) was found, compared to the healthy control group (~3 %). Moreover, 6.3 % of patients carried two *CFTR* variants, making CFTR-RD a possibility in these patients and thus strengthening the argument that AWP can be considered a CFTR-RD.

7.3. CFTR-RD AWP diagnostic criteria and recommendations for follow-up and treatment

CF should be excluded in individuals presenting with AWP, particularly if symptoms in other systems co-exist, regardless of their severity [121,124]. If the CF diagnostic criteria are not satisfied then CFTR-RD could still be the diagnosis so further testing should be considered, particularly if the symptoms are painful and unresponsive to standard treatment (see below). Previous studies have not confirmed a relationship between sweat chloride concentration and the severity of AWP and data on other functional tests in AWP (e.g. nasal PD and ICM) are not available [115,118]. Therefore, it is recommended that genetic analysis is the first diagnostic test, particularly as AWP is frequently found in individuals without any CFTR involvement. When a single *CFTR* variant is found, further assessment is recommended. The genotypes that have been reported in AWP are the same as those found in the other conditions associated with CFTR-RD.

7.4. Alternative causes of AWP and how to exclude them

Other conditions associated with AWP include marasmus, nephrotic syndrome, cardiac abnormalities and drug reactions [114,117, 125–127]. Genetic causes, other than *CFTR*, have been indentified, including heterozygosity for variants in the *Aquaporin 5* (*AQP5*) gene in 3 % of 198 French patients with isolated AWP [128]. *AQP5* variants have been described in autosomal-dominant diffuse Non-Epidermolytic

Palmoplantar Keratoderma of Bothnian type, which is a permanent condition of wrinkling, worsened by water immersion [129]. Heterozygosity for *CFTR* and *AQP5* variants could increase the risk of AWP, as previously suggested [130]. The inclusion of *CFTR* and *AQP5* genes in the genetic exploration of AWP is thus worth considering.

7.5. Recommendations for follow-up and treatment of CFTR-RD AWP

Patients presenting with AWP and at least one *CFTR* variant, should be referred to a CF center for sweat testing and a complete clinical evaluation with appropriate follow-up [1]. Referral for genetic counselling is recommended.

Usually specific treatment is not needed. If AWP interferes with daily life, local topical aluminium hydroxide should be considered for first line treatment. Botulinum toxin injection has been described as an alternative treatment in some cases [131].

8. Conclusions

This is the third paper of the ECFS CFTR-RD project, covering the individual disorders, including their clinical features, route to diagnosis, alternative causes and key elements to their management. By undertaking this process, it became clear early on, that the evidence underpinning each disorder, is highly variable. This was not only the situation for the ‘established’ disorders (i.e. CAVD, bronchiectasis and pancreatitis) but particularly for the ‘emerging’ pathologies. Although this was not unexpected, it presented challenges reaching definitive conclusions for some conditions, which explains the variable presentation of the information and recommendations. Table 5 summarises the paper, providing an overview for the reader, with areas of uncertainty highlighted. Notwithstanding this, the paper provides an important up-to-date overview of the conditions associated with CFTR dysfunction and, importantly, extends into new areas, explaining the principles of their diagnosis based on *CFTR* genetics and electrophysiological testing, and the exclusion of other (often more common) causes. However, as

each condition has its own unique set of clinical, practical and cost challenges, it is acknowledged that unified recommendations across all of them is not realistic at the current time, although with greater recognition of CFTR-RD across the specialities from this programme, we hope this can be achieved in the future. It is clear, that gaps in our knowledge remain, including an accurate estimate of the prevalence of each condition, the precise level of CFTR dysfunction and why some organs are affected more than others. Producing evidence showing that the accurate diagnosis of these conditions leads to a change in treatment and long-term outcomes, and the specific role of CFTR modulators, is an area in urgent need of further research. This, along with the role of registries, genetic counselling and effective dissemination of CFTR-RD knowledge beyond the CF community, are the topics in the next and final paper of this project.

Funding

No funding received.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

NJS reports consulting fees and/or honoraria from Vertex, Chiesi, Gilead and Menarini. KDB reports consulting fees and/or honoraria from Vertex, ELoxx, Splisense and Arcturus. JGM reports honoraria from Vertex, Chiesi, Abbvie and Viartis. PGM reports consulting fees and/or honoraria from Vertex, AstraZenica, Limbic and MedEd. CS reports consulting fees and/or honoraria from Vertex, Chiesi, TFF and Pfizer. JC reports consulting fees for Astrazeneca, Boehringer Ingelheim, Grifols, Gilead sciences, Insmmed, Genentech, Glaxosmithkline, Antabio, Zambon and Trudell. OCY reports consulting fees and honoraria from Vertex. DGD reports consulting fees and/or honoraria from Chiesi, Insmmed and Vertex. DP reports payments for educational talks. DVD reports consulting fees for Arcturus, Arrevus, BiomX, Clarametyx, CFF, Enbiotix,

Table 5  
Summary of the disorders considered a CFTR-RD and key diagnostic, genetic and clinical features.

|                                                        | CAVD                                                                                                                                                  | Diffuse Bronchiectasis                                                                                                   | Chronic or acute recurrent pancreatitis                                     | Chronic rhinosinusitis                                           | ABPA                                                                               | Primary Sclerosing Cholangitis                 | Aquagenic wrinkling of the palms                                                                                        |
|--------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------|------------------------------------------------------------------|------------------------------------------------------------------------------------|------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------|
| <b>Diagnostic confirmation*</b>                        | Semen analysis<br>Clinical examination<br>Imaging (US)                                                                                                | High resolution thoracic CT                                                                                              | Serum amylase/lipase<br>MRCP                                                | Clinical examination<br>Nasendoscopy<br>CT<br>±MRI<br>Allergic   | Aspergillus serology (total IgE, specific IgE, IgG)<br>±skin prick tests<br>Asthma | MRCP<br>ERCP                                   | Clinical examination<br>Biopsy – not usually required                                                                   |
| <b>Main alternative diagnoses/causes to consider**</b> | Renal pathology                                                                                                                                       | Immunological, Autoimmune, Post infective                                                                                | Gallstones, Alcohol, Toxic/metabolic causes                                 |                                                                  |                                                                                    | Autoimmune                                     | Marasmus, nephrotic syndrome, cardiac abnormalities, drug reactions                                                     |
| <b>Other genes implicated</b>                          | <i>ADGRG2</i>                                                                                                                                         | PCD genes<br><i>ENaC</i>                                                                                                 | <i>SPINK1</i><br><i>PRSS1</i><br><i>CTRC</i><br><i>TRPV6</i><br><i>CPA1</i> |                                                                  | <i>PLA2G4</i> ,<br><i>PLA2G2D</i><br><i>ALOX15</i>                                 |                                                | <i>AQP5</i>                                                                                                             |
| <b>Uncertainties and research questions</b>            | The role of MRI.<br>The extent of disease – bilateral/unilateral disease or partial obstruction.<br>The role of CFTR modulators and disease reversal. | The role of other non-CFTR genes.<br>The effectiveness of CF-specific treatment, e.g. dornase alpha and CFTR modulators. | The impact of non-CFTR genes.<br>The role of CFTR modulators.               | The accuracy of nasal PD in CRS.<br>The role of CFTR modulators. | The role of CFTR modulators.                                                       | The mechanistic significance of CFTR function. | The mechanistic significance of CFTR function.<br>The true prevalence in the general population, CF and other CFTR-RDs. |

Key: \*the principal modalities/tests for reaching the diagnosis of the individual condition; \*\*not exhaustive (see the main text for a more complete list); US=ultrasound; CT=computed tomography; ERCP=endoscopic retrograde cholangiopancreatography; MRCP=magnetic resonance cholangiopancreatography; MRI=magnetic resonance imaging; PD=potential difference.

Microbion, Pyros, Respirion and Zambon. ISG reports honoraria from Vertex. KWS, FB, VV, EDW, MW, EB, DD, PE, EG, GH, AK, DN, JAN, CR, IS, MDW report no personal fees/honoraria

## Acknowledgements

The authors wish to acknowledge and thank all members of the ECFS Diagnostic Network Working Group- for their support, comments, discussion and contributions throughout this project.

## References

- Castellani C, De Boeck K, De Wachter E, et al. ECFS standards of care on CFTR-related disorders: updated diagnostic criteria. *J Cyst Fibros* 2022;21:908–21.
- Sermet-Gaudelus I, Girodon E, Vermeulen F, et al. ECFS standards of care on CFTR-related disorders: diagnostic criteria of CFTR dysfunction. *J Cyst Fibros* 2022;21:922–36.
- Weiske WH, Salzler N, Schroeder-Printzen I, et al. Clinical findings in congenital absence of the vasa deferentia. *Andrologia* 2000;32:13–8.
- Baydilli N, Gokce A, Karabulut SY, et al. Klinefelter's syndrome with unilateral absence of vas deferens. *Fertil Steril* 2010;94: 1529 e1521–1529 e1522.
- Fedder J, Cruger D, Oestergaard B, et al. Etiology of azoospermia in 100 consecutive nonvasectomized men. *Fertil Steril* 2004;82:1463–5.
- Oates RD, Amos JA. The genetic basis of congenital bilateral absence of the vas deferens and cystic fibrosis. *J Androl* 1994;15:1–8.
- Bieth E, Hamdi SM, Miesusset R. Genetics of the congenital absence of the vas deferens. *Hum Genet* 2021;140:59–76.
- Daudin M, Bieth E, Bujan L, et al. Congenital bilateral absence of the vas deferens: clinical characteristics, biological parameters, cystic fibrosis transmembrane conductance regulator gene mutations, and implications for genetic counseling. *Fertil Steril* 2000;74:1164–74.
- Cai Z, Li H. Congenital bilateral absence of the vas deferens. *Front Genet* 2022;13: 775123.
- Li L, Liang C. Ultrasonography in diagnosis of congenital absence of the vas deferens. *Med Sci Monit* 2016;22:2643–7.
- Akinsal EC, Baydilli N, Dogan ME, et al. Comorbidity of the congenital absence of the vas deferens. *Andrologia* 2018.
- Schlegel PN, Shin D, Goldstein M. Urogenital anomalies in men with congenital absence of the vas deferens. *J Urol* 1996;155:1644–8.
- Minhas S, Bettocchi C, Boeri L, et al. European association of urology guidelines on male sexual and reproductive health: 2021 update on male infertility. *Eur Urol* 2021;80:603–20.
- Chiang HS, Lin YH, Wu YN, et al. Advantages of magnetic resonance imaging (MRI) of the seminal vesicles and intra-abdominal vas deferens in patients with congenital absence of the vas deferens. *Urology* 2013;82:345–51.
- Patat O, Pagin A, Siegfried A, et al. Truncating mutations in the adhesion G protein-coupled receptor G2 gene ADGRG2 cause an X-linked congenital bilateral absence of vas deferens. *Am J Hum Genet* 2016;99:437–42.
- Cai H, Qing X, Niringiyumukiza JD, et al. CFTR variants and renal abnormalities in males with congenital unilateral absence of the vas deferens (CUAVD): a systematic review and meta-analysis of observational studies. *Genet Med* 2019; 21:826–36.
- Teng H, Jorissen M, Van Poppel H, et al. Increased proportion of exon 9 alternatively spliced CFTR transcripts in vas deferens compared with nasal epithelial cells. *Hum Mol Genet* 1997;6:85–90.
- Castellani C, Cuppens H, Macek Jr M, et al. Consensus on the use and interpretation of cystic fibrosis mutation analysis in clinical practice. *J Cyst Fibros* 2008;7:179–96.
- Miesusset R, Bieth E, Daudin M, et al. Male partners of infertile couples with congenital unilateral absence of the vas deferens are mainly non-azoospermic. *Andrology* 2020;8:645–53.
- Kolettis PN, Sandlow JL. Clinical and genetic features of patients with congenital unilateral absence of the vas deferens. *Urology* 2002;60:1073–6.
- Bush A, Floto RA. Pathophysiology, causes and genetics of paediatric and adult bronchiectasis. *Respirology* 2019;24:1053–62.
- Kapur N, Masel JP, Watson D, et al. Bronchoarterial ratio on high-resolution CT scan of the chest in children without pulmonary pathology: need to redefine bronchial dilatation. *Chest* 2011;139:1445–50.
- Matsuoka S, Uchiyama K, Shima H, et al. Bronchoarterial ratio and bronchial wall thickness on high-resolution CT in asymptomatic subjects: correlation with age and smoking. *AJR Am J Roentgenol* 2003;180:513–8.
- Polverino E, Goeminne PC, McDonnell MJ, et al. European respiratory society guidelines for the management of adult bronchiectasis. *Eur Respir J* 2017;50.
- Hill AT, Sullivan AL, Chalmers JD, et al. British Thoracic Society guideline for bronchiectasis in adults. *Thorax* 2019;74:1–69.
- Pignatti PF, Bombieri C, Benetazzo M, et al. CFTR gene variant IVS8-5T in disseminated bronchiectasis. *Am J Hum Genet* 1996;58:889–92.
- Girodon E, Cazeneuve C, Leborgy F, et al. CFTR gene mutations in adults with disseminated bronchiectasis. *Eur J Hum Genet* 1997;5:149–55.
- Bombieri C, Benetazzo M, Saccomani A, et al. Complete mutational screening of the CFTR gene in 120 patients with pulmonary disease. *Hum Genet* 1998;103: 718–22.
- King PT, Freezer NJ, Holmes PW, et al. Role of CFTR mutations in adult bronchiectasis. *Thorax* 2004;59:357–8.
- Colak Y, Nordestgaard BG, Afzal S. Morbidity and mortality in carriers of the cystic fibrosis mutation CFTR Phe508del in the general population. *Eur Respir J* 2020;56.
- Puechal X, Fajac I, Bienvenu T, et al. Increased frequency of cystic fibrosis deltaF508 mutation in bronchiectasis associated with rheumatoid arthritis. *Eur Respir J* 1999;13:1281–7.
- Agarwal R, Khan A, Aggarwal AN, et al. Link between CFTR mutations and ABPA: a systematic review and meta-analysis. *Mycoses* 2012;55:357–65.
- Lawless D, Allen HL, Thaventhiran JED, et al. Prevalence of CFTR variants in primary immunodeficiency patients with bronchiectasis is an important modifying cofactor. *J Allergy Clin Immunol* 2023;152:257–65.
- Cartier Y, Kavanagh PV, Johkoh T, et al. Bronchiectasis: accuracy of high-resolution CT in the differentiation of specific diseases. *AJR Am J Roentgenol* 1999;173:47–52.
- Lynch DA, Newell J, Hale V, et al. Correlation of CT findings with clinical evaluations in 261 patients with symptomatic bronchiectasis. *AJR Am J Roentgenol* 1999;173:53–8.
- Reiff DB, Wells AU, Carr DH, et al. CT findings in bronchiectasis: limited value in distinguishing between idiopathic and specific types. *AJR Am J Roentgenol* 1995; 165:261–7.
- Hubert D, Fajac I, Bienvenu T, et al. Diagnosis of cystic fibrosis in adults with diffuse bronchiectasis. *J Cyst Fibros* 2004;3:15–22.
- Nick JA, Chacon CS, Brayshaw SJ, et al. Effects of gender and age at diagnosis on disease progression in long-term survivors of cystic fibrosis. *Am J Respir Crit Care Med* 2010;182:614–26.
- Behan L, Dimitrov BD, Kuehni CE, et al. PICADAR: a diagnostic predictive tool for primary ciliary dyskinesia. *Eur Respir J* 2016;47:1103–12.
- Lucas JS, Barbato A, Collins SA, et al. European respiratory society guidelines for the diagnosis of primary ciliary dyskinesia. *Eur Respir J* 2017;49.
- Smyth AR, Bell SC, Bojcin S, et al. European cystic fibrosis society standards of care: best practice guidelines. *J Cyst Fibros* 2014;13(Suppl 1):S23–42.
- Castellani C, Duff AJA, Bell SC, et al. ECFS best practice guidelines: the 2018 revision. *J Cyst Fibros* 2018;17:153–78.
- Whitcomb DC, Pancreatitis North American Pancreatitis Study G. TIGAR-O version 2 risk/etiology checklist with topic reviews, updates, and use primers. *Clin Transl Gastroenterol* 2019;10:e00027.
- Garipey CE, Heyman MB, Lowe ME, et al. Causal evaluation of acute recurrent and chronic pancreatitis in children: consensus from the INSPIRE group. *J Pediatr Gastroenterol Nutr* 2017;64:95–103.
- Masson E, Chen JM, Cooper DN, et al. PRSS1 copy number variants and promoter polymorphisms in pancreatitis: common pathogenetic mechanism, different genetic effects. *Gut* 2018;67:592–3.
- Bishop MD, Freedman SD, Zielinski J, et al. The cystic fibrosis transmembrane conductance regulator gene and ion channel function in patients with idiopathic pancreatitis. *Hum Genet* 2005;118:372–81.
- Mayerle J, Sendler M, Hegyi E, Genetics et al. Cell biology, and pathophysiology of pancreatitis. *Gastroenterology* 2019;156:1951–68. e1951.
- Kumar S, Ooi CY, Werlin S, et al. Risk factors associated with pediatric acute recurrent and chronic pancreatitis: lessons from INSPIRE. *JAMA Pediatr* 2016; 170:562–9.
- Uc A, Husain SZ. Pancreatitis in children. *Gastroenterology* 2019;156:1969–78.
- LaRusch J, Jung J, General LJ, et al. Mechanisms of CFTR functional variants that impair regulated bicarbonate permeation and increase risk for pancreatitis but not for cystic fibrosis. *PLoS Genet* 2014;10:e1004376.
- Angyal D, Bijvelds MJC, Bruno MJ, et al. Bicarbonate transport in cystic fibrosis and pancreatitis. *Cells* 2021;11.
- Ramsey ML, Li SS, Lara LF, et al. Cystic fibrosis transmembrane conductance regulator modulators and the exocrine pancreas: a scoping review. *J Cyst Fibros* 2023;22:193–200.
- McKay IR, Ooi CY. The exocrine pancreas in cystic fibrosis in the era of CFTR modulation: a mini review. *Front Pediatr* 2022;10:914790.
- Akshintala VS, Kamal A, Faghii M, et al. Cystic fibrosis transmembrane conductance regulator modulators reduce the risk of recurrent acute pancreatitis among adult patients with pancreas sufficient cystic fibrosis. *Pancreatology* 2019; 19:1023–6.
- Ramsey ML, Gokun Y, Sobotka LA, et al. Cystic fibrosis transmembrane conductance regulator modulator use is associated with reduced pancreatitis hospitalizations in patients with cystic fibrosis. *Am J Gastroenterol* 2021;116: 2446–54.
- Gould MJ, Smith H, Rayment JH, et al. CFTR modulators increase risk of acute pancreatitis in pancreatic insufficient patients with cystic fibrosis. *J Cyst Fibros* 2022;21:600–2.
- Lee DJ, Sykes J, Griffin K, et al. The negative impact of chronic rhinosinusitis on the health-related quality of life among adult patients with cystic fibrosis. *J Cyst Fibros* 2022;21:800–6.
- Wang X, Kim J, McWilliams R, et al. Increased prevalence of chronic rhinosinusitis in carriers of a cystic fibrosis mutation. *Arch Otolaryngol Head Neck Surg* 2005;131:237–40.
- Raman V, Clary R, Siegrist KL, et al. Increased prevalence of mutations in the cystic fibrosis transmembrane conductance regulator in children with chronic rhinosinusitis. *Pediatrics* 2002;109:E13.
- Fokkens WJ, Lund VJ, Hopkins C, et al. Executive summary of EPOS 2020 including integrated care pathways. *Rhinology* 2020;58:82–111.

- [61] Eggesbo HB, Dolvik S, Stiris M, et al. Complementary role of MR imaging of ethmoidmaxillary sinus disease depicted at CT in cystic fibrosis. *Acta Radiol* 2001; 42:144–50.
- [62] Schraven SP, Wehrmann M, Wagner W, et al. Prevalence and histopathology of chronic polypoid sinusitis in pediatric patients with cystic fibrosis. *J Cyst Fibros* 2011;10:181–6.
- [63] Sommerburg O, Wielputz MO, Trame JP, et al. Magnetic resonance imaging detects chronic rhinosinusitis in infants and preschool children with cystic fibrosis. *Ann Am Thorac Soc* 2020;17:714–23.
- [64] Beswick DM, Humphries SM, Balkissoon CD, et al. Olfactory dysfunction in cystic fibrosis: impact of CFTR modulator therapy. *J Cyst Fibros* 2021.
- [65] Stapleton AL, Kimple AJ, Goralski JL, et al. Elexacaftor-Tezacaftor- Ivacaftor improves sinonasal outcomes in cystic fibrosis. *J Cyst Fibros* 2022;21:792–9.
- [66] Mainz JG, Arnold C, Wittstock K, et al. Ivacaftor reduces inflammatory mediators in upper airway lining fluid from cystic fibrosis patients with a G551D mutation: serial non-invasive home-based collection of upper airway lining fluid. *Front Immunol* 2021;12:642180.
- [67] Pranke IM, Hatton A, Simonin J, et al. Correction of CFTR function in nasal epithelial cells from cystic fibrosis patients predicts improvement of respiratory function by CFTR modulators. *Sci Rep* 2017;7:7375.
- [68] Wang X, Moylan B, Leopold DA, et al. Mutation in the gene responsible for cystic fibrosis and predisposition to chronic rhinosinusitis in the general population. *JAMA* 2000;284:1814–9.
- [69] Moller ME, Alanin MC, Gronhøj C, et al. Sinus bacteriology in patients with cystic fibrosis or primary ciliary dyskinesia: a systematic review. *Am J Rhinol Allergy* 2017;31:293–8.
- [70] Hatton A, Bergougnot A, Zybert K, et al. Reclassifying inconclusive diagnosis after newborn screening for cystic fibrosis. Moving forward. *J Cyst Fibros* 2022; 21:448–55.
- [71] Lindig J, Steger C, Beiersdorf N, et al. Smell in cystic fibrosis. *Eur Arch Otorhinolaryngol* 2013;270:915–21.
- [72] Hentschel J, Muller U, Dohr F, et al. Influences of nasal lavage collection-, processing- and storage methods on inflammatory markers—evaluation of a method for non-invasive sampling of epithelial lining fluid in cystic fibrosis and other respiratory diseases. *J. Immunol. Methods* 2014;404:41–51.
- [73] Mainz JG, Nährlich L, Schien M, et al. Concordant genotype of upper and lower airways P aeruginosa and S aureus isolates in cystic fibrosis. *Thorax* 2009;64: 535–40.
- [74] Arndal E, Johansen HK, Haagenen JAJ, et al. Primary ciliary dyskinesia patients have the same P. Aeruginosa clone in sinuses and lungs. *Eur Respir J* 2020;55.
- [75] Thamboo A, Santos RC, Naidoo L, et al. Use of the SNOT-22 and UPSIT to appropriately select pediatric patients with cystic fibrosis who should be referred to an otolaryngologist: cross-sectional study. *JAMA Otolaryngol Head Neck Surg* 2014;140:934–9.
- [76] Spielman DB, Beswick DM, Kimple AJ, et al. The management of cystic fibrosis chronic rhinosinusitis: an evidenced-based review with recommendations. *Int Forum Allergy Rhinol* 2022;12:1148–83.
- [77] Beer H, Southern KW, Swift AC. Topical nasal steroids for treating nasal polyposis in people with cystic fibrosis. *Cochrane Database Syst Rev* 2011;CD008253.
- [78] Möller W, Schuschnig U, Celik G, et al. Topical drug delivery in chronic rhinosinusitis patients before and after sinus surgery using pulsating aerosols. *PLoS ONE* 2013;8:e74991.
- [79] Mainz JG, Schädlich K, Schien C, et al. Sinonasal inhalation of tobramycin vibrating aerosol in cystic fibrosis patients with upper airway Pseudomonas aeruginosa colonization: results of a randomized, double-blind, placebo-controlled pilot study. *Drug Des Devel Ther* 2014;8:209–17.
- [80] Mainz JG, Schien C, Schiller I, et al. Sinonasal inhalation of dornase alfa administered by vibrating aerosol to cystic fibrosis patients: a double-blind placebo-controlled cross-over trial. *J Cyst Fibros* 2014;13:461–70.
- [81] Aanaes K, Alanin MC, Nielsen KG, et al. The accessibility of topical treatment in the paranasal sinuses on operated cystic fibrosis patients assessed by scintigraphy. *Rhinology* 2018;56:268–73.
- [82] Mainz JG, Koitschev A. Pathogenesis and management of nasal polyposis in cystic fibrosis. *Curr Allergy Asthma Rep* 2012;12:163–74.
- [83] DiMango E, Overdevest J, Keating C, et al. Effect of highly effective modulator treatment on sinonasal symptoms in cystic fibrosis. *J Cyst Fibros* 2021;20:460–3.
- [84] Wucherpfennig L, Triphan SMF, Wege S, et al. Magnetic resonance imaging detects improvements of pulmonary and paranasal sinus abnormalities in response to elexacaftor/tezacaftor/ivacaftor therapy in adults with cystic fibrosis. *J Cyst Fibros* 2022;21:1053–60.
- [85] Ramos KJ, Guimbellot JS, Valapour M, et al. Use of elexacaftor/tezacaftor/ivacaftor among cystic fibrosis lung transplant recipients. *J Cyst Fibros* 2022;21: 745–52.
- [86] Middleton PG, Chen SC, Meyer W. Fungal infections and treatment in cystic fibrosis. *Curr Opin Pulm Med* 2013;19:670–5.
- [87] Agarwal R, Chakrabarti A, Shah A, et al. Allergic bronchopulmonary aspergillosis: review of literature and proposal of new diagnostic and classification criteria. *Clin Exp Allergy* 2013;43:850–73.
- [88] Stevens DA, Moss RB, Kurup VP, et al. Allergic bronchopulmonary aspergillosis in cystic fibrosis—state of the art: cystic fibrosis foundation consensus conference. *Clin Infect Dis* 2003;37(Suppl 3):S225–64.
- [89] Mastella G, Rainisio M, Harms HK, et al. Allergic bronchopulmonary aspergillosis in cystic fibrosis. A European epidemiological study. Epidemiologic registry of cystic fibrosis. *Eur Respir J* 2000;16:464–71.
- [90] Geller DE, Kaplowitz H, Light MJ, et al. Allergic bronchopulmonary aspergillosis in cystic fibrosis: reported prevalence, regional distribution, and patient characteristics. scientific advisory group, investigators, and coordinators of the epidemiologic study of cystic fibrosis. *Chest* 1999;116:639–46.
- [91] Gamaletsos MN, Hayes G, Harris C, et al. F508del CFTR gene mutation in patients with allergic bronchopulmonary aspergillosis. *J Asthma* 2018;55:837–43.
- [92] Marchand E, Verellen-Dumoulin C, Mairesse M, et al. Frequency of cystic fibrosis transmembrane conductance regulator gene mutations and 5T allele in patients with allergic bronchopulmonary aspergillosis. *Chest* 2001;119:762–7.
- [93] Miller PW, Hamosh A, Macek Jr M, et al. Cystic fibrosis transmembrane conductance regulator (CFTR) gene mutations in allergic bronchopulmonary aspergillosis. *Am J Hum Genet* 1996;59:45–51.
- [94] Bickford JS, Mueller C, Newsom KJ, et al. Effect of allergy and inflammation on eicosanoid gene expression in CFTR deficiency. *J Cyst Fibros* 2013;12:258–65.
- [95] Muller C, Braag SA, Herlihy JD, et al. Enhanced IgE allergic response to aspergillus fumigatus in CFTR-/- mice. *Lab Invest* 2006;86:130–40.
- [96] Mueller C, Torrez D, Braag S, et al. Partial correction of the CFTR-dependent ABPA mouse model with recombinant adeno-associated virus gene transfer of truncated CFTR gene. *J Gene Med* 2008;10:51–60.
- [97] Antunes J, Fernandes A, Borrego LM, et al. Cystic fibrosis, atopy, asthma and ABPA. *Allergol Immunopathol (Madr)* 2010;38:278–84.
- [98] Patel AR, Patel AR, Singh S, et al. Treating allergic bronchopulmonary aspergillosis: a review. *Cureus* 2019;11:e4538.
- [99] Chetty A. Pathology of allergic bronchopulmonary aspergillosis. *Front Biosci* 2003;8:e110–4.
- [100] Pashley CH. Fungal culture and sensitisation in asthma, cystic fibrosis and chronic obstructive pulmonary disorder: what does it tell us? *Mycopathologia* 2014;178: 457–63.
- [101] Sunman B, Ademhan Tural D, Ozsezen B, et al. Current approach in the diagnosis and management of allergic bronchopulmonary aspergillosis in children with cystic fibrosis. *Front Pediatr* 2020;8:582964.
- [102] Durieu I, Pellet O, Simonot L, et al. Sclerosing cholangitis in adults with cystic fibrosis: a magnetic resonance cholangiographic prospective study. *J Hepatol* 1999;30:1052–6.
- [103] Strandvik B, Hjelte L, Gabrielson N, et al. Sclerosing cholangitis in cystic fibrosis. *Scand J Gastroenterol Suppl* 1988;143:121–4.
- [104] Girodon E, Sternberg D, Chazouilleres O, et al. Cystic fibrosis transmembrane conductance regulator (CFTR) gene defects in patients with primary sclerosing cholangitis. *J Hepatol* 2002;37:192–7.
- [105] Gallegos-Orozco JF, EY C, Wang N, et al. Lack of association of common cystic fibrosis transmembrane conductance regulator gene mutations with primary sclerosing cholangitis. *Am J Gastroenterol* 2005;100:874–8.
- [106] Sheth S, Shea JC, Bishop MD, et al. Increased prevalence of CFTR mutations and variants and decreased chloride secretion in primary sclerosing cholangitis. *Hum Genet* 2003;113:286–92.
- [107] Pall H, Zielenski J, Jonas MM, et al. Primary sclerosing cholangitis in childhood is associated with abnormalities in cystic fibrosis-mediated chloride channel function. *J Pediatr* 2007;151:255–9.
- [108] Henckaerts L, Jaspers M, Van Steenberghe W, et al. Cystic fibrosis transmembrane conductance regulator gene polymorphisms in patients with primary sclerosing cholangitis. *J Hepatol* 2009;50:150–7.
- [109] Werlin S, Scotet V, Uguen K, et al. Primary sclerosing cholangitis is associated with abnormalities in CFTR. *J Cyst Fibros* 2018;17:666–71.
- [110] Di Giorgio A, Vergani D, Mieli-Vergani G. Cutting edge issues in juvenile sclerosing cholangitis. *Dig Liver Dis* 2022;54:417–27.
- [111] Lalwani N, Bhargava P, Chintapalli KN, et al. Current update on primary and secondary sclerosing cholangitis. *Curr Probl Diagn Radiol* 2011;40:248–61.
- [112] Elliott RB. Letter: skin wrinkling in cystic fibrosis. *Lancet* 1974;2:1383.
- [113] English 3rd JC, McCollough ML. Transient reactive papulotranslucent acrokeratoderma. *J Am Acad Dermatol* 1996;34:686–7.
- [114] Atzori L, Ferrelli C, Rongioletti F. Aquagenic (pseudo) keratoderma (aquagenic palmo-plantar keratoderma, aquagenic wrinkling of palms). *Clin Dermatol* 2021; 39:256–60.
- [115] Arkin LM, Flory JH, Shin DB, et al. High prevalence of aquagenic wrinkling of the palms in patients with cystic fibrosis and association with measurable increases in transepidermal water loss. *Pediatr Dermatol* 2012;29:560–6.
- [116] Alexopoulos A, Choularas G, Kakourou T, et al. Aquagenic wrinkling of the palms after brief immersion to water test as a screening tool for cystic fibrosis diagnosis. *J Eur Acad Dermatol Venereol* 2021;35:1717–24.
- [117] Paul GR, Bai S, Jackson K, et al. Aquagenic wrinkling in children under two years of age: could this be a potential clinical referral tool for cystic fibrosis among non-screened populations? *J Cyst Fibros* 2021;20:e77–83.
- [118] Berk DR, Ciliberto HM, Sweet SC, et al. Aquagenic wrinkling of the palms in cystic fibrosis: comparison with controls and genotype-phenotype correlations. *Arch Dermatol* 2009;145:1296–9.
- [119] Rongioletti F, Tomasini C, Crovato F, et al. Aquagenic (pseudo) keratoderma: a clinical series with new pathological insights. *Br J Dermatol* 2012;167:575–82.
- [120] Weil B, Chaillou E, Troussier F, et al. [Aquagenic palmo-plantar keratoderma in children with cystic fibrosis]. *Arch Pediatr* 2013;20:1306–9.
- [121] Torres-Laboy PM, Melendez-Montanez JM, De Jesus-Rojas W. Aquagenic wrinkling of the palm: a rare diagnostic clue of cystic fibrosis and the response to CFTR-modulating therapy. *Cureus* 2021;13:e14425.
- [122] Grasmann H, Ratjen F, Solomon M. Aquagenic wrinkling of the palms in a patient with cystic fibrosis. *N Engl J Med* 2013;369:2362–3.
- [123] Raynal C, Girodon E, Audrezet MP, et al. CFTR gene variants: a predisposition factor to aquagenic palmo-plantar keratoderma. *Br J Dermatol* 2019;181:1097–9.
- [124] Stewart LC, Doe SJ, Bourke SJ, et al. Aquagenic palmar wrinkling as a presenting feature of cystic fibrosis gene dysfunction. *Clin Exp Dermatol* 2009;34:e647–9.

- [125] Bouwman K, Menichino S, Kruithof I, et al. Two new cases of aquagenic wrinkling of the palms and literature review on drug interactions. *Dermatol Online J* 2020; 26.
- [126] Gualdi G. Aquagenic wrinkling: the importance to be known. *J Eur Acad Dermatol Venereol* 2016;30:e224–5.
- [127] Wilder-Smith EP. Aquagenic wrinkling of the palms is due to vasoconstriction of palmar skin vasculature. *Med Hypotheses* 2013;81:963–5.
- [128] Sperelakis-Beedham B, Lopez M, Bourrat E, et al. AQP5, a second gene at play with CFTR in aquagenic palmoplantar keratoderma. *J Eur Acad Dermatol Venereol* 2023;37:e654–7.
- [129] Blaydon DC, Lind LK, Plagnol V, et al. Mutations in AQP5, encoding a water-channel protein, cause autosomal-dominant diffuse nonepidermolytic palmoplantar keratoderma. *Am J Hum Genet* 2013;93:330–5.
- [130] Castellani C, Boner AL. Aquagenic wrinkling and cystic fibrosis carriership: a dubious relationship. *Eur J Intern Med* 2018;57:e43.
- [131] Yang K, Zhou C, Luke J. Aquagenic wrinkling of the palms: review of the literature. *Clin Exp Dermatol* 2022;47:1910–5.